

Cours on Lie groups

(fall 2018)

Below you find an incomplete list of DEFINITIONS, *examples* and *theorems* from the course on Lie groups:

REVIEW: TOPOLOGICAL MANIFOLDS, SMOOTH MANIFOLDS, SMOOTH MAPS, SUBMANIFOLDS, REGULAR VALUE

pre-image of a regular value is a submanifold (without proof)

LIE GROUPS, HOMOMORPHISM BETWEEN LIE GROUPS

G a Lie group, $H \subset G$ a subgroup which is also a submanifold $\implies H$ is a Lie group

$S^1, T^n, Gl(n, \mathbb{R}), Gl(n, \mathbb{C}), O(n), U(n), SO(n), SU(n)$, *symplectic group* $Sp(n)$, *dimension of these Lie groups*

REVIEW: TANGENT VECTORS, DERIVATIONS, TANGENT SPACE, TANGENT BUNDLE, DIFFERENTIAL OF A SMOOTH MAP, VECTOR FIELDS

LEFT INVARIANT VECTOR FIELDS

Lie group is parallizable, Lie bracket $[X, Y]$ of two left invariant vector fields is left invariant, Jacobi identity

LIE ALGEBRA $(LG, [,])$, INTEGRAL CURVES, ONE-PARAMETER GROUPS, EXPONENTIAL MAP $\exp : LG \rightarrow G$

$(\exp_*)_0 = id_{LG}$, naturality of the exponential map, \exp for a matrix group $G \subset Gl(n, \mathbb{C})$

Lie algebras of classical groups, $sl(n, \mathbb{R}), o(n), so(n), u(n), su(n), sp(n)$

G connected \implies a homomorphism $f : G \rightarrow H$ is determined by $f_* : LG \rightarrow LH$. G a Lie group and H a closed abstract subgroup $\implies H$ is a submanifold (and, hence, H a Lie (sub)group).

G, H Lie groups and $f : G \rightarrow H$ continuous homomorphism of groups $\implies f$ is smooth (and, hence, a homomorphism of Lie groups).

ADJOINT $\text{Ad} : G \rightarrow \text{Aut}(LG)$, $\text{ad} : LG \rightarrow \text{End}(LG)$

$[X, Y] = \text{ad}(X)(Y)$. G abelian $\implies [,] = 0$.

For G connected holds: G abelian $\iff \exp : LG \rightarrow G$ is surjective.

G abelian connected $\implies G \cong T^k \times \mathbb{R}^l$.

G compact abelian $\implies G \cong T^n \times B$ for B a finite abelian group.

COMPLEX G -REPRESENTATION $\rho : G \times V \rightarrow V$, REPRESENTATION SPACE V , ASSOCIATED HOMOMORPHISM $G \rightarrow \text{Aut}(V)$, MATRIX REPRESENTATION, MORPHISM BETWEEN G -REPRESENTATIONS, ISOMORPHIC/EQUIVALENT REPRESENTATIONS, UNITARY REPRESENTATIONS

standard $U(n)$ -representation \mathbb{C}^n , one-dimensional representations of S^1 and of T^n , adjoint representation Ad , direct sum of representations, tensor product of representations, dual representation, conjugate representation, exterior algebra, symmetric algebra, representations $\Lambda^k V$, $S^k V$

SUBREPRESENTATION, IRREDUCIBLE REPRESENTATION, COMPLETELY REDUCIBLE REPRESENTATION

Schur's lemma. G abelian and V irreducible G -representation $\implies \dim V = 1$.

G compact \implies there exists a left invariant normalized integral $\int : C^0(G) \rightarrow \mathbb{R}$, every G -representation has a G -invariant scalar product and every G -representation is completely reducible.

CHARACTER, IRREDUCIBLE CHARACTER, CLASS FUNCTION

properties of the character: 2.19 and theorem 2.20

character of S^1 - and T^n -representations

$\text{Irr}(G; \mathbb{C})$, $d_W : \text{Hom}_G(V, W) \otimes W \rightarrow V$, MULTIPLICITY FOR AN IRREDUCIBLE REPRESENTATION IN A GIVEN REPRESENTATION

$d : \bigoplus_{W \in \text{Irr}(G; \mathbb{C})} \text{Hom}_G(V, W) \otimes W \rightarrow V$ is an isomorphism of G -representations.

From now on all groups are assumed to be compact!

The isomorphism type of a G -representation V is uniquely determined by its character.

$\langle \chi_V, \chi_V \rangle = 1 \implies V$ is irreducible.

V irreducible G -representation, W irreducible H -representation $\implies V \otimes W$

irreducible $(G \times H)$ -representation.

Every $(G \times H)$ -representation is sum of representations of the form $V \otimes W$, where V is an irreducible G -representation and W is an irreducible H -representation.

representations of $SU(2)$ and $SO(3)$

$$Irr(SU(2); \mathbb{C}) = \{V_0, V_1, V_2, \dots\}, Irr(SO(3); \mathbb{C}) = \{V_0, V_2, V_4, \dots\}$$

Clebsch-Gordan formulas

GROTHENDIECK CONSTRUCTION, COMPLEX REPRESENTATION RING $R(G)$

$$R(SU(2)) \cong \mathbb{Z}[V_1]$$

$$G \text{ and } H \text{ compact Lie groups} \implies R(G \times H) \cong R(G) \otimes R(H).$$

$$R(T^r), R(SU(2) \times S^1)$$

orthogonality relations 3.1

REPRESENTATIVE FUNCTIONS, $\mathcal{T}(G; \mathbb{C})$

$\mathcal{T}(G; \mathbb{C})$ is a subalgebra of $C^0(G; \mathbb{C})$ and $\overline{\mathcal{T}(G; \mathbb{C})} = \mathcal{T}(G; \mathbb{C})$

$$\mathcal{T}(S^1; \mathbb{C})$$

(Peter-Weyl) G compact $\implies \mathcal{T}(G; \mathbb{C})$ dense in $(L^2(G), \|\cdot\|)$.

$\mathcal{T}(G; \mathbb{C})$ dense in $(C^0(G; \mathbb{C}), \|\cdot\|_1)$.

The irreducible characters generate a dense subspace in the vector space of all continuous class functions (wrt. $\|\cdot\|_1$).

Every compact Lie group G admits a faithful representation.

Every compact Lie group G is isomorphic to a closed subgroup of $U(N)$ for some $N \gg 0$.

MAXIMAL TORUS T OF G , NORMALIZER $N_G(T)$, WEYL GROUP $W(G) := N_G(T)/T$

maximal tori exist

Suppose $H \subset G$ is a subgroup. Then: H is a maximal torus $\iff H$ is a maximal connected abelian subgroup,

$N_G(T)_e = T$ and $W(G)$ is a finite group.

Suppose G is connected and T is a max. torus. Then: every element is conjugate to an element in T (proof uses the Lefschetz fixed point formula),

all maximal tori are conjugate to each other.

For G connected the exponential map is surjective.

For G connected, S a connected abelian subgroup of G and $g \in Z_G(S)$ exists a maximal torus which contains S and g .

RANK $\text{rk}(G)$ OF A COMPACT LIE GROUP

Suppose G is connected, \tilde{T} is a maximal torus and S is a connected abelian subgroup. Then: $Z_G(\tilde{T}) = \tilde{T}$, $Z_G(S)$ is the union of all max. tori which contain S ,

the center $Z(G)$ is the intersection of all max. tori.

For G connected the Weyl group acts effectively on T ,

two elements $x, y \in T$ are conjugate in $G \iff \exists w \in W(G)$ with $w(x) = y$.

Suppose G is connected. Then: $R(G) \rightarrow R(T)^{W(G)}$ is injective and $R(G)$ is isomorphic to a subring of $\mathbb{Z}[\lambda_1, \lambda_1^{-1}, \dots, \lambda_r, \lambda_r^{-1}]$, where $r := \text{rk}(G)$.

maximal tori, Weyl group and representation ring for $U(n)$...